



Active Risk Management Program in Operating Rooms of Hospitals in Different Countries: A Scoping Review Study

Akbar Javan Biparva ¹, Raana Gholamzadeh Nikjoo ^{1,*}, Ali Jannati ¹, Mohamad Arab ² and Ali Ostadi ³

¹Department of Health Policy and Health Services Management, School of Management and Medical Informatics, Iranian Center of Excellence in Health Management, Tabriz Health Services Management Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

²Department of Management and Health Economic, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

³Clinical research Development Unit, Sina Hospital, Tabriz University of Medical Sciences, Tabriz, Iran

*Corresponding author: Department of Health Policy and Health Services Management, School of Management and Medical Informatics, Iranian Center of Excellence in Health Management, Tabriz Health Services Management Research Center, Tabriz University of Medical Sciences, Tabriz, Iran. Email: r.gholamzade@gmail.com

Received 2022 December 24; Revised 2023 March 13; Accepted 2023 April 16.

Abstract

Context: Risk management in hospitals is essential for improving healthcare quality. This study analyzed the characteristics of risk management programs implemented in operating rooms of hospitals across countries worldwide.

Evidence Acquisition: This study was a scoping review of online database studies, including Web of Knowledge, PubMed, Scopus, Cochrane, Springer, ProQuest, Iranian SID, and Magiran databases and the Google Scholar search engine. Three people independently performed the study selection, quality assessment, data extraction, and analysis among studies that reviewed risk management programs in health systems and those conducted outside the health system. Articles in non-English languages (including Persian) were excluded.

Results: A total of 21 studies with similar purposes and data collection methods were included. The characteristics of risk management programs were classified into six main factors: Objectives, components, steps, results, prerequisites, facilitators of risk management programs, and 35 sub-factors.

Conclusions: The conceptual framework of any risk management program should include at least the objectives: Risk eradication, safety promotion, quality improvement and prevention and reduction of risks, component: Communication and monitoring; steps: (1) preoperative evaluation during (logging); (2) evaluation during surgery; (3) post-surgery evaluation (logout); and the results: Achieving effective methods in reducing errors; prerequisites: Human resource, knowledge and information, and facilitators such as the use of monitoring technologies and error detection and reporting in the operating room.

Keywords: Risk, Risk Management, Operating Room

1. Context

Nowadays, developing strategies to improve the quality of care plays a vital role in international health policies. This factor has prompted governments to improve the quality of care. The four components of quality, from the perspective of the World Health Organization, which are the basis for the formation of clinical service support are professional performance (technical quality), resource use (productivity), patient satisfaction with the services provided, and risk management (risk of injury or illness in connection with the services provided) (1).

Risk management in health care refers to a diverse

group of measures performed to improve the quality and ensure the safety of services for patients. The daily concern of high-risk organizations, including hospitals, is the management of unexpected events (2). Risk management is crucial in improving health care quality, effective communication between hospital staff and patients, patient desirability, and Limiting abortion practices in hospitals (3, 4). In the past, risk management in clinical settings was examined with a reactive view. In other words, risk management was performed after the incident, and its causes and factors were analyzed to prevent its recurrence. However, nowadays, the emphasis is on the active method in which risk probability is accepted and managed adequately before the threat occurs (5).

Studies showed that the operating room is only a small part of the total hospital in terms of physical space, comprising less than 4% of the total hospital space and less than 3% of the total hospital staff, but 35% of the complaints related to risks related to the operating room (6). Other studies reported that the risk rate in the operating room is 71.8% and more than 50 unknown types (7). Therefore, risk management in the operating room should be prioritized over other hospital sections. About 234 million surgeries are performed annually worldwide (8, 9). Using a systematic and comprehensive method in surgical departments to prevent the possibility of error will bring good results. However, even in industrialized countries, error prevention is not done properly (10). Surgery is an important healthcare service accounting for 40% of hospital costs. Also, due to the higher probability of error in the operating room and its consequences for patients, the performance of this department is very effective in assessing the quality of services provided to patients and their satisfaction (11, 12). The high error rate in intensive care units, operating rooms (ORs), and emergency departments might be explained by the complex nature of the operation (13). Among the above-mentioned healthcare facilities, operating rooms are more concerned with high-risk operations. Therefore, errors are more likely for patients and staff in operating rooms (14). These errors are classified into eight general themes: Operational risks, technology, legal, human capacity, financial, strategic, clinical, and patient safety and risk (15).

Errors and adverse events occur due to inefficient management and other technical, human, and organizational inadequacies (e.g., lack of proper communication and training and insufficient standard methods). These cases have led to issues such as wrong place surgery and wrong anesthesia management, which are significant in operating rooms and endanger patients' lives and impose huge costs on healthcare systems (14). Among other incidents in the operating room, we can mention the possibility of physical damage due to electric shock, burns, fire, contact with blood products, and inhalation of toxic substances. Some devices used in the operating room, such as lasers, radiography equipment, and chemical sterilizers, can also lead to long-term damage if the staff is careless in the security field (16). Therefore, some effective factors in the occurrence of medical errors in the operating room can be prevented, reducing errors and adverse operating room accidents and leading to increased quality of care (14).

2. New Contributions

Numerous studies have used different models for risk management in the operating room. In Guo's study, for risk management in the operating room, a risk quantification matrix and a risk registration form were implemented to identify potential hazards, and then policies were designed and implemented to reduce or eliminate these hazards (17). A counseling mechanism and risk monitoring system were also used to minimize risks among operating room nurses (17). In a study by DeRosier et al. on operating room risk management, a surgical risk assessment model was implemented using failure modes and effects analysis (FMEA), and multi-objective optimization based on analysis was presented (18). This work aimed to determine the most important risk points during the surgical process and analyze the relevant solutions (18). So far, no study has attempted to provide a comprehensive picture of the characteristics of risk management programs implemented in different countries. Therefore, this study aimed to implement a scoping review approach to analyze and review the characteristics of risk management programs implemented in the operating room of hospitals in various countries worldwide. Our results will provide a comprehensive picture of this program to help develop a risk management program in the operating room of hospitals.

3. Evidence Acquisition

In this study, we used Arksey and O'Malley's framework (19). The framework includes six steps: Identifying the research question, identifying relevant studies, selecting/screening studies, scaling/segmenting data, summarizing, reporting results, and optional consultation with experts on findings. The detailed explanation of the six main steps is as follows:

(1) Identifying the research question: Like systematic review studies, in this method, the starting point is identifying the research question on which search strategies are built. The study question is: What are the characteristics of risk management programs implemented in the operating room of hospitals in various countries worldwide?

(2) Identification of related studies: The scoping review focuses on identifying the original studies (published and gray studies) and appropriate review studies to answer the main research question as comprehensively as possible. In the present study, appropriate keywords and their combination, such as proactive risk management (PRM), proactive risk assessment, risk management, health

systems, hospitals, operating room, objectives, steps, components, outcomes, facilitators, and prerequisites, were searched in various databases such as Web of Knowledge, PubMed, Scopus, Cochrane, Springer, ProQuest, Iranian databases, including SID, Magiran, and Google search engine. All published articles and reports in this field were selected. The search was done on December 21, 2022. In addition to manually searching the journals and sources of selected articles, organizational reports, published government documents, websites, and other available information and gray resources were also reviewed. Based on the search, about 11,030 related articles were obtained (Figure 1).

(3) Selection/screening of studies: All stages of selection and screening of articles were performed separately by two research team members. Disputes were resolved through discussion, and if necessary, they were referred to a third party with more information and experience. First, the titles of all studies reviewed and articles inconsistent with the study's objectives were removed. In the following steps, the abstract and full text of the articles were reviewed to exclude studies that met exclusion criteria and had poor relevance to study objectives. This study was performed without a time limit and with the following inclusion and exclusion criteria. Inclusion criteria included studies that reviewed the risk management program in health systems. Exclusion criteria included studies conducted outside the health system and non-English and Persian articles. Out of 11,030 articles in this study, about 5,175 were removed due to duplication. Also, 5,715 articles were removed due to the mismatch of the title and abstract of the article with the main purpose. Likewise, about 119 articles were removed due to the mismatch of the article's main text with the main objective; finally, 21 articles were extracted (Figure 1).

(4) Data segmentation: This stage arranges key items from the reviewed reports. The results of the selected studies were managed through a data extraction form designed by the research team, including the authors, year of publication, type of risk, purpose, steps, and components of risk management. Two people independently extracted the information from the selected articles, and the ambiguities were resolved in consultation with other research team members.

(5) Summarizing and reporting the results: The results were analyzed using thematic analysis. This way, a preliminary study was conducted to identify the extracted studies. As a result, a thematic framework based on codes specific to the studies was designed. Another researcher reviewed each study, and the findings were extracted in specific codes. In the next step, the main themes were extracted following the semantic framework and

connection and sometimes integration between codes, and the findings were organized within the main themes. Finally, after ensuring the validity of the main themes, they were organized based on their fit with the main research question of the category and then presented in the form of a table of findings.

(6) Presenting practical tips and recommendations: After performing the previous steps, based on the extracted data and the opinions of the research team members, tips and recommendations were made in the form of article discussion and knowledge translation.

4. Results

Based on the presented framework, six main themes of findings were extracted, including objectives, components, steps, results, prerequisites, and facilitators of the risk management program. The goals of the risk management program included seven sub-themes: Quality improvement, safety improvement, cost reduction, satisfaction increase, risk prevention and reduction, financial loss protection, and risk eradication. Among these goals are risk eradication with 62%, safety improvement with 57%, quality improvement and prevention and reduction of risks with 52%, protection of financial losses with 14%, reduction of costs with 10%, and increase of satisfaction with 5%, which are the most frequent purposes among the articles (Table 1).

The components of the risk management program include six sub-themes: Resources, leadership, organizational culture, communication, monitoring, and training. Among these components, communication (57%), supervision (52%), resources (43%), leadership (33%), organizational culture (24%), and education (19%) were the most frequent risk management components among the articles (Table 2).

The stages of the risk management program include four main models; model 3 (57%) and models 1, 2, and 4 (14%) were the most frequent risk management steps among the articles (Table 3). The results of the risk management program include four sub-themes consisting of identifying the main causes of errors, achieving effective methods in reporting errors, achieving effective methods in reducing errors, and the need for informed consent. Achieving effective methods in reducing errors (57%), achieving effective methods in reporting errors (38%), identifying the main causes of errors (24%), and the need for informed consent (5%) were the most common types of the risk management program among the articles (Table 4).

Prerequisites for the risk management program include five sub-themes: Human resources, financial




Figure 1. PRISMA diagram for selected studies

resources, culture and structure, knowledge and information, and equipment and technology. Among these prerequisites, human resources (100%), knowledge and information (71%), culture and structure (24%), equipment and technology (19%), and financial resources (10%) were the most prevalent types of prerequisites in the risk management program across the articles (Table 5).

Facilitators of the risk management program include nine sub-themes: Use of modeling, use of expert opinions, teamwork, use of quality approach, use of equipment with good performance and adequate supervision, presence of a person as a risk manager, use of records, use of monitoring technologies to identify and report faults and identifying and monitoring safety indicators. Among the facilitators, the use of monitoring and detection technologies (57%), teamwork (38%), the use of equipment with good performance and adequate supervision (14%), the use of expert opinions (10%), the use of modeling, using a qualitative approach, and using records and identifying and monitoring safety indicators (5%) were most frequent in risk management program facilitators

among the articles (Table 6).

5. Discussion

The present study attempted to identify the characteristics of risk management programs in the operating rooms of hospitals in different countries. A total of 21 studies that were the same in terms of purpose and data collection methods were included in the study. After analyzing the findings, the characteristics of risk management programs were classified into six main factors and 35 sub-factors.

The highest number of studies on risk management in the operating room was in the United States (42%). The oldest study was conducted in 1980, and the most recent one was performed in 2019. Eleven articles have mentioned all the errors that can happen in the operating room. Six of the articles are related to clinical errors, and two of the articles are related to fire in the operating room. Also, one of the articles is related to human errors, and one is related to equipment failure. Most articles have used the

Table 2. Components of Risk Management Programs in the Operating Rooms of Hospitals in Different Countries

No.	Author Name and Year of Publication	Components of the Risk Management Program					
		References	Leadership	Organizational Culture	Connections	Monitoring	Education
1	Sayed et al., 2013 (20)	*				*	
2	Guo, 2015 (17)				*	*	
3	van Beuzekom et al., 2012 (21)	*					*
4	Schimpff, 2007 (22)	*	*	*	*		*
5	Borie et al., 2018 (23)	*				*	
6	Frabetti et al., 2005 (24)	*				*	
7	Amghar et al., 2017 (25)		*				
8	Silen-Lipponen et al., 2005 (26)	*	*				
9	Pomerantz et al., 2011 (27)					*	*
10	DeRosier et al., 2019 (18)		*		*	*	
11	Kavosi et al., 2017 (28)		*		*	*	
12	Le Bourg and Gallois, 2011 (29)			*	*		
13	Lepanluoma et al., 2013 (30)			*	*		
14	American College of Obstetricians and Gynecologists, 2010 (31)	*			*	*	
15	American Society of Anesthesiologists Task Force on Operating Room Fires et al., 2008 (32)				*		
16	Bower, 2002 (33)			*	*		
17	McLain, 1980 (34)	*	*		*		*
18	Alfredsdottir and Bjornsdottir, 2008 (35)		*	*	*		
19	Tabibzadeh and Jahangiri, 2018 (14)					*	
20	Guedon et al., 2014 (36)				*	*	
21	Frosini et al., 2016 (37)	*				*	

FMEA, root-cause analysis (RCA) model, and WHO safety guidelines as risk management models.

One of the important features of risk management programs in hospitals is the goals of these programs. Many hospitals plan and implement risk management programs to address the risks involved. The existence of an incident registration system is an active approach to risk management, which is opposed to a passive approach. In the proactive approach, their repetition can be prevented by recording incidents and learning from mistakes (38, 39). The next main goal is to improve safety in the operating room. There is a strong link between risk management and

safety promotion (40). In an organization with a positive safety culture, all employees at different levels value safety and consider it the organization's main priority (41). In such an organization, the likelihood of dangerous events and even healthcare costs are significantly reduced (42). Attention to prospective risk management approaches will be very effective in increasing safety in the hospital (43). Quality improvement has also been one of the goals of the risk management program in the operating room (40). Due to the increasing competition in the field of medicine, one of the approaches of hospital managers to increase the quality of care is to apply the risk management

Table 4. Risk Management Program Results in the Operating Rooms of Hospitals in Different Countries

No.	Author Name and Year of Publication	Risk Management Program Results			
		Identifying the Main Causes of Errors	Achieving Effective Methods in Reporting Errors	Achieving Effective Methods to Reduce Errors	The Need for Informed Consent
1	Sayed et al., 2013 (20)	*		*	*
2	Guo, 2015 (17)			*	
3	van Beuzekom et al., 2012 (21)		*	*	
4	Schimpff, 2007 (22)			*	
5	Borie et al., 2018 (23)	*			
6	Frabetti et al., 2005 (24)		*		
7	Amghar et al., 2017 (25)		*		
8	Silen-Lipponen et al., 2005 (26)		*		
9	Pomerantz et al., 2011 (27)			*	
10	DeRosier et al., 2019 (18)			*	
11	Kavosi et al., 2017 (28)	*			
12	Le Bourg and Gallois, 2011 (29)			*	
13	Lepanluoma et al., 2013 (30)	*	*		
14	American College of Obstetricians and Gynecologists, 2010 (31)			*	
15	American Society of Anesthesiologists Task Force on Operating Room Fires et al., 2008 (32)			*	
16	Bower, 2002 (33)	*			
17	McLain, 1980 (34)		*	*	
18	Alfredsdottir and Bjornsdottir, 2008 (35)	*			
19	Tabibzadeh and Jahangiri, 2018 (14)			*	
20	Guedon et al., 2014 (36)		*		
21	Frosini et al., 2016 (37)		*		

process comprehensively and efficiently (44). So, in all quality improvement programs, the risk management approach is one of the main axes of creating, deploying, and using management systems in organizations (45). Risk prevention and reduction are also among the main goals of risk management programs in the operating room. In general, risk management means preventing the occurrence of the hazard, reducing risks and injuries, and managing risks after an event (44, 46). Therefore, by implementing this program, hospitals tend to prevent risks and reduce their error rate. The next goal is to protect

against financial losses and reduce operating room costs. Hospitals can take steps to reduce clinical error-related loss by using a risk management approach (44). The ultimate goal of implementing a risk management program is to improve satisfaction. According to studies in this field, risk management, in turn, will lead to improved satisfaction (46).

The next feature of operating room risk management programs is the program's components. What are the components of risk management programs designed and implemented for hospital operating rooms? The first

Table 5. Prerequisites for Risk Management Programs in the Operating Rooms of Hospitals in Different Countries

No.	Author Name and Year of Publication	Prerequisites for Risk Management Program				
		Human Resources	Financial Resources	Culture and Structure	Knowledge and Information	Equipment and Technology
1	Sayed et al., 2013 (20)	*			*	
2	Guo, 2015 (17)	*			*	
3	van Beuzekom et al., 2012 (21)	*	*			
4	Schimpff, 2007 (22)	*		*		*
5	Borie et al., 2018 (23)	*				
6	Frabetti et al., 2005 (24)	*				*
7	Amghar et al., 2017 (25)	*			*	
8	Silen-Lipponen et al., 2005 (26)	*			*	
9	Pomerantz et al., 2011 (27)	*			*	
10	DeRosier et al., 2019 (18)	*			*	
11	Kavosi et al., 2017 (28)	*			*	
12	Le Bourg and Gallois, 2011 (29)	*		*	*	
13	Lepanluoma et al., 2013 (30)	*			*	
14	American College of Obstetricians and Gynecologists, 2010 (31)	*			*	
15	American Society of Anesthesiologists Task Force on Operating Room Fires et al., 2008 (32)	*			*	
16	Bower, 2002 (33)	*	*	*	*	
17	McLain, 1980 (34)	*		*	*	
18	Alfredsdottir and Bjornsdottir, 2008 (35)	*		*	*	
19	Tabibzadeh and Jahangiri, 2018 (14)	*			*	
20	Guedon et al., 2014 (36)	*				*
21	Frosini et al., 2016 (37)	*				*

and most important component of communication risk management programs is communication. Effective risk management programs require open lines of communication (47). According to studies, increased cooperation between operating room staff and staff communication has reduced patient risks and mortality

(30). To carry out the risk management program in the operating room correctly, it is necessary to form a team and communicate effectively between the teams to identify and manage the risks through these communications. The next important component of the risk management program is monitoring. Until the monitoring is done,

Table 6. Risk Management Program Facilitators in the Operating Rooms of Hospitals in Different Countries

No.	Author Name and Year of Publication	Risk Management Program Facilitator								
		Use of Modeling	Use of the Opinions of Experts	Teamwork	Use of a Qualitative Approach	Having an Individual as a Risk Manager	Use of Records and Documents	Monitoring, Detection, and Error Reporting Technologies	Identifying and Monitoring Safety Indicators	Equipment with Good Performance and Adequate Supervision
1	Sayed et al., 2013 (20)				*		*			
2	Guo, 2015 (17)							*		
3	van Beuzekom et al., 2012 (21)							*		
4	Schimpff, 2007 (22)							*		
5	Borie et al., 2018 (23)							*		
6	Frabetti et al., 2005 (24)							*		
7	Amghar et al., 2017 (25)	*								
8	Silen-Lipponen et al., 2005 (26)		*	*				*		
9	Pomerantz et al., 2011 (27)							*		
10	DeRosier et al., 2019 (18)			*						*
11	Kavosi et al., 2017 (28)			*				*		
12	Le Bourg and Gallois, 2011 (29)			*				*		
13	Lepanluoma et al., 2013 (30)							*		
14	American College of Obstetricians and Gynecologists, 2010 (31)			*						*
15	American Society of Anesthesiologists Task Force on Operating Room Fires et al., 2008 (32)		*	*						
16	Bower, 2002 (33)			*		*				
17	McLain, 1980 (34)					*		*		
18	Alfredsdottir and Bjornsdottir, 2008 (35)			*						
19	Tabilzadeh and Jahangiri, 2018 (14)								*	
20	Guedon et al., 2014 (36)									*
21	Frosini et al., 2016 (37)							*		

it will not be determined how well the program will work. Risk and emergencies can be reduced by applying and monitoring clear preventive policies with specific frameworks and roles in the operating room (30). Therefore, establishing hospital monitoring programs to implement a risk management program is necessary (48). Resources are also a component of risk management programs. Managing human, financial, material, support, and system resources is critical for each resource before,

during, and after an unforeseen event. Undesirable hidden risk is designed to help improve patient safety in the operating room (49). The next pillar of the risk management program is leadership. Leaders in every hospital must be aware of the critical need for patient safety programs and strive to reduce risks, and make safety champions of change (50). Encouraging commitment in employees to implement the program, the teamwork of members, execution of orders, information transfer, case

reporting, etc., align with the concept of leadership (51).

Organizational culture is also a component of the risk management program. As explained above, successful organizations implementing risk management programs have a positive safety culture, and all employees at different levels value safety and consider it the organization's main priority (41). Studies in this field consider a culture of prevention and protection against hazards by nursing staff in the operating room to be crucial in increasing safety (52). The last component of the risk management program is training. Training is one of the human resources approaches requiring extensive internal training for properly implementing the error reporting system and error management steps in the operating room (47).

The next important feature is the implementation of the risk management program in the hospital. In general, four models for risk management in the operating room were identified, the most common of which are related to the steps: (1) preoperative evaluation during (logging); (2) evaluation during surgery; (3) post-surgery evaluation (logout). In this model, which can also be called pre-test/post-test evaluation (49), using different methods, the hospital first identifies the risks before surgery and then performs its proposed interventions during surgery to identify errors. Finally, after surgery, the effectiveness of the measures is re-evaluated. Other models of risk management (creating an environment, identifying, analyzing, evaluating, risk elimination, consulting, monitoring, creating and informing prevention programs, creating a channel for error reporting, training, and obtaining feedback) among the steps used in the operating room risk management have been reported to reduce the risks after implementing a risk management program in the operating room.

The next feature of risk management programs is the results of the program. What are the results of the implemented risk management programs for the hospital or the operating room? According to the findings, the highest results obtained from implementing risk management programs were the achievement of effective methods in reducing errors. For example, the use of a safe surgical checklist (46), the use of well-functioning equipment under adequate supervision (31), and systematic reporting to assist in decision-making in the areas of risk management (53) were among the effective methods in reducing errors. The next important result of implementing a risk management program in the operating room is achieving effective methods for reporting errors. A safety notification system, which is a system for monitoring the safety status (periodic repairs and recorded failures) of operating room devices and

facilitating information about failures (36), a web-based monitoring dashboard (37), and automatic collection of computer data using technology (54) were among the methods of reporting errors. Identifying the main causes of errors is also one of the main results of implementing a risk management program. In a study, 204 errors were reported for 36 sub-processes, with the highest frequency related to human and organizational errors and the lowest to technical errors (28). In another study, nurses identified the most common causes of the error, including lack of knowledge, information, and supervision, heavy workload, and poor judgment, which may lead to active and hidden errors (52). Another study identified the necessity of obtaining informed consent. The importance of completing and clarifying surgical satisfaction for the patient or one of the patient's relatives, which is done by increasing the level of communication between surgeons and patients and discussing the conditions and possible results of surgeries, will lead the hospital to be safe against possible financial and legal consequences and risks (55).

The next feature of risk management programs is the prerequisites of the risk management program. Prerequisites are a set of factors that are required to run the program. Human resources are the most important prerequisite for implementing a hospital risk management program. In the absence of human resources, no management program will be feasible because the implementers of this program are human resources. In most articles, discussing human resources and teamwork in the operating room is recognized as a necessity of risk management programs (29, 30). The next prerequisite is the discussion of knowledge and information. Some sources believe that risk management in an organization cannot be successfully implemented without knowledge and information (56). Leaders and officials need the necessary knowledge to deal with the crisis at the right time and place. The next prerequisite is culture and structure. In most articles, culture and structure are the basic principles for implementing a risk management program (51, 57). Unless there is a clear structure for staff in a hospital and the staff does not follow a certain culture, one cannot expect to implement a risk management program in the operating room. The next issue is equipment and technology. In risk management, using technologies and equipment, especially in the communication and information for organizations involved in operations, is very important. Collecting data from the risk area using these technologies will be much faster and more reliable and help better implement the risk management program, for example, the automatic collection of computer data using technology to monitor infections in the operating room (54), the use of a safety

status system to monitor the safety of operating room equipment (36), and use of a monitoring dashboard to collect information from devices in the room action (37). Financial resources are the last prerequisite for implementing risk management programs in the operating room. The implementation of any program requires financial resources to be operational. In implementing a risk management program in the operating room, the provision of financial resources such as human resources and knowledge is of particular importance (49).

Risk management program facilitators are the final feature of operating room risk management programs. Facilitators are tools that facilitate the implementation of these programs. Among the facilitators, the use of monitoring technologies and error detection and reporting are recognized as the most important types of facilitators of risk management program implementation. These technologies could be prerequisites for any program, making receiving errors easier and taking timely action. The next facilitator is teamwork. Operating room teamwork means combining complementary skills, overlapping team members, and mutual understanding while caring for patients undergoing surgery (58). Evidence has shown that teamwork in healthcare systems improves patient safety (59). For this reason, most studies have used the workforce required in the risk management program as a team. The next case is using equipment with good performance and adequate supervision, which facilitates control and reduces the occurrence of hazards in the operating room (31). Using experts' opinions also facilitates risk management in the operating room; for example, advice based on a combination of scientific literature and analysis of expert opinions to facilitate patient care in the face of risks (31). Subsequent facilitators include modeling that provides a framework for presenting causal relationships and enables possible inference among a set of variables (25). It is necessary to act with reflection and commitment at all times with the aim of permanent improvement (29), the use of records and documents such as medical history or any particular medical condition (55), and identifying and monitoring safety indicators that are used to assess and monitor internal and external social and technical factors in a health care setting (16).

5.1. Conclusions

Based on the results of the current study, we identified six main factors as features of the risk management program in the operating room. The operators of the risk management program in the operating room should pay special attention to these factors during

planning in order to get the most effectiveness from the implementation of the program. The conceptual framework of any risk management program should include at least the objectives: Risk eradication, safety promotion, quality improvement and prevention and reduction of risks, component: communication and monitoring; steps: (1) preoperative evaluation during (logging); (2) evaluation during surgery; (3) post-surgery evaluation (logout); and the results: Achieving effective methods in reducing errors, prerequisites: Human resource, knowledge and information, and facilitators such as the use of monitoring technologies and error detection and reporting in the operating room. Using this framework, any risk management program operator can adapt their conditions to the program objectives, the components that the program must have, the steps it must go through, the prerequisites and facilitators, and the results it wants to achieve, and, as a result, take the appropriate route to reduce hazards in the operating room.

These results help health insurance organizations and health policymakers get information about the risk management frameworks and the status of the operating rooms of the hospitals to carry out managerial and policy interventions for the effective implementation of the developed framework. The results will also help the health system achieve its set goals to improve people's health, meet their reasonable expectations, and receive safe and line-free service for patients and a safe work environment for personnel.

5.2. Limitations

The main limitation of this research was conducting very limited studies in the field of risk management, especially in the operating room of hospitals worldwide. However, we tried to deal with this limitation using organizational reports, published government documents, websites, and other gray information sources for obtaining information.

Footnotes

Authors' Contribution: A. J. B., R. Gh., and A. J. conceptualized the study and participated in the initial study design. M. A. and A. O. obtained the research funding. All authors reviewed the manuscript. A. J. B. coordinated manuscript revisions. The authors read and approved the final manuscript.

Conflict of Interests: The authors declare no conflict of interest.

Funding/Support: The authors received no financial support for the research, authorship, and/or publication of this article.

References

- Mitra A. *Fundamentals of quality control and improvement*. Hoboken, USA: John Wiley & Sons; 2016.
- Amaral T, Costa APC. Improving decision-making and management of hospital resources: An application of the PROMETHEE II method in an Emergency Department. *Oper Res Health Care*. 2014;3(1):1-6. <https://doi.org/10.1016/j.orhc.2013.10.002>.
- Braithwaite J, Wears RL, Hollnagel E. Resilient health care: turning patient safety on its head. *Int J Qual Health Care*. 2015;27(5):418-20. [PubMed ID: 26294709]. <https://doi.org/10.1093/intqhc/mzv063>.
- Aita A, Padoan A, Antonelli G, Sciacovelli L, Plebani M. Patient safety and risk management in medical laboratories: theory and practical application. *J Lab Precis Med*. 2017;2:75. <https://doi.org/10.21037/jlpm.2017.08.14>.
- Farokhzadian J, Dehghan Nayeri N, Borhani F. Assessment of Clinical Risk Management System in Hospitals: An Approach for Quality Improvement. *Glob J Health Sci*. 2015;7(5):294-303. [PubMed ID: 26156927]. [PubMed Central ID: PMC4803839]. <https://doi.org/10.5539/gjhs.v7n5p294>.
- Stones J, Yates D. Clinical risk assessment tools in anaesthesia. *BJA Educ*. 2019;19(2):47-53. [PubMed ID: 33456869]. [PubMed Central ID: PMC7807823]. <https://doi.org/10.1016/j.bjae.2018.09.009>.
- Fraser JRS, Simkins BJ. The challenges of and solutions for implementing enterprise risk management. *Bus Horiz*. 2016;59(6):689-98. <https://doi.org/10.1016/j.bushor.2016.06.007>.
- Nitassi S, Belayachi J, Chihab M, Rkaini I, Benayad J, Benbouzid MA, et al. Evaluation of Post Laryngectomy Pharyngocutaneous Fistula risk Factors. *Iran J Otorhinolaryngol*. 2016;28(85):141-7. [PubMed ID: 27280101]. [PubMed Central ID: PMC4881883].
- Burcharth J, Falkenberg A, Schack A, Ekeloef S, Gogenur I. The effects of early enteral nutrition on mortality after major emergency abdominal surgery: A systematic review and meta-analysis with Trial Sequential Analysis. *Clin Nutr*. 2021;40(4):1604-12. [PubMed ID: 33744604]. <https://doi.org/10.1016/j.clnu.2021.02.050>.
- Rah JE, Manger RP, Yock AD, Kim GY. A comparison of two prospective risk analysis methods: Traditional FMEA and a modified healthcare FMEA. *Med Phys*. 2016;43(12):6347. [PubMed ID: 27908165]. <https://doi.org/10.1118/1.4966129>.
- Lai H, Gemming L. Approaches to patient satisfaction measurement of the healthcare food services: A systematic review. *Clin Nutr ESPEN*. 2021;42:61-72. [PubMed ID: 33745623]. <https://doi.org/10.1016/j.clnesp.2020.12.019>.
- Ahmadi A, Mosallanezhad B. Using fuzzy FMEA to increase patient safety in fundamental processes of operating room. *Journal of Industrial and Systems Engineering*. 2018;11(3):146-66.
- Bates DW, Singh H. Two Decades Since To Err Is Human: An Assessment Of Progress And Emerging Priorities In Patient Safety. *Health Aff (Millwood)*. 2018;37(11):1736-43. [PubMed ID: 30395508]. <https://doi.org/10.1377/hlthaff.2018.0738>.
- Tabibzadeh M, Jahangiri G. A Proactive Risk Assessment Framework to Enhance Patient Safety in Operating Rooms. *Los Angeles, USA*. 2018. p. 263-74.
- Carroll RL, Hoppes M, Hagg-Rickert S, Youngberg BJ, McCarthy BA, Shope D, et al. *Enterprise risk management: A framework for success*. Chicago, USA: American Society For Healthcare Risk Management; 2014.
- Zheng W, Shuai J, Shan K. The energy source based job safety analysis and application in the project. *Saf Sci*. 2017;93:9-15. <https://doi.org/10.1016/j.ssci.2016.11.009>.
- Guo L. Implementation of a risk management plan in a hospital operating room. *Int J Nurs Sci*. 2015;2(4):348-54. <https://doi.org/10.1016/j.ijnss.2015.10.007>.
- DeRosier JM, Hansemann BK, Smith-Wheelock MW, Bagian JP. Using Proactive Risk Assessment (HFMEA) to Improve Patient Safety and Quality Associated with Intraocular Lens Selection and Implantation in Cataract Surgery. *Jt Comm J Qual Patient Saf*. 2019;45(10):680-5. [PubMed ID: 31422905]. <https://doi.org/10.1016/j.jcjq.2019.06.003>.
- Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol*. 2005;8(1):19-32. <https://doi.org/10.1080/1364557032000119616>.
- Sayed HA, Zayed M, El Qareh NM, Khafagy H, Helmy AH, Soliman M. Patient safety in the operating room at a governmental hospital. *J Egypt Public Health Assoc*. 2013;88(2):85-9. [PubMed ID: 23963087]. <https://doi.org/10.1097/01.EPX.0000430955.28520.e5>.
- van Beuzekom M, Boer F, Akerboom S, Hudson P. Patient safety in the operating room: an intervention study on latent risk factors. *BMC Surg*. 2012;12:10. [PubMed ID: 22726757]. [PubMed Central ID: PMC3407486]. <https://doi.org/10.1186/1471-2482-12-10>.
- Schimpff SC. Improving operating room and perioperative safety: background and specific recommendations. *Surg Innov*. 2007;14(2):127-35. [PubMed ID: 17558019]. <https://doi.org/10.1177/1553350607301746>.
- Borie F, Mathonnet M, Deleuze A, Millat B, Gravie JF, Johanet H, et al. Risk management for surgical energy-driven devices used in the operating room. *J Visc Surg*. 2018;155(4):259-64. [PubMed ID: 29289460]. <https://doi.org/10.1016/j.jvisurg.2017.12.003>.
- Frabetti A, Vandini A, Pantoja Rodriguez S, Margelli F, Cavicchioli M, Migliori M, et al. Microbiological risk in operating rooms: new strategies for infections surveillance. *Milan, Italy*. 2005. p. 275-86.
- Amghar M, Zoullouti B, Sbiti N. Risk Analysis of Operating Room Using the Fuzzy Bayesian Network Model. *International Journal of Engineering*. 2017;30(1). <https://doi.org/10.5829/idosi.ije.2017.30.01a.09>.
- Silen-Lipponen M, Tossavainen K, Turunen H, Smith A. Potential errors and their prevention in operating room teamwork as experienced by Finnish, British and American nurses. *Int J Nurs Pract*. 2005;11(1):21-32. [PubMed ID: 15610341]. <https://doi.org/10.1111/j.1440-172X.2005.00494.x>.
- Pomerantz RG, Lee DA, Siegel DM. Risk assessment in surgical patients: balancing iatrogenic risks and benefits. *Clin Dermatol*. 2011;29(6):669-77. [PubMed ID: 22014989]. <https://doi.org/10.1016/j.clindermatol.2011.08.012>.
- Kavosi Z, Setoodehzadeh F, Fardid M, Gholami M, Khojastefar M, Hatam M, et al. [Risk assessment of the processes of operating room department using the failure mode and effects analysis (FMEA) method]. *Journal of Hospital*. 2017;16(3):57-70. Persian.
- Le Bourg M, Gallois PM. Managing risk in the operating room: from theory to practice. *Orthop Traumatol Surg Res*. 2011;97(4 Suppl):S49-54. [PubMed ID: 21531649]. <https://doi.org/10.1016/j.otsr.2011.03.008>.
- Lepanluoma M, Takala R, Kotkansalo A, Rahi M, Ikonen TS. Surgical safety checklist is associated with improved operating room safety culture, reduced wound complications, and unplanned readmissions in a pilot study in neurosurgery. *Scand J Surg*. 2014;103(1):66-72. [PubMed ID: 24345978]. <https://doi.org/10.1177/1457496913482255>.
- American College of Obstetricians and Gynecologists. *Patient safety in the surgical environment. Committee Opinion No. 464. American College of Obstetricians and Gynecologists*. Washington, USA: American College of Obstetricians and Gynecologists; 2010.
- Caplan RA, Barker SJ, Connis RT, Cowles C, de Richemond AL; American Society of Anesthesiologists Task Force on Operating Room Fires, et al. Practice advisory for the prevention and management of operating room fires. *Anesthesiology*. 2008;108(5):786-801. quiz 971-2. [PubMed ID: 18431114]. <https://doi.org/10.1097/01.anes.0000299343.87119.a9>.

33. Bower JO. Designing and implementing a patient safety program for the OR. *AORN J.* 2002;**76**(3):452-6. [PubMed ID: [12227288](#)]. [https://doi.org/10.1016/s0001-2092\(06\)61659-1](https://doi.org/10.1016/s0001-2092(06)61659-1).
34. McLain NB. Risk management in the operating room. *AORN J.* 1980;**31**(5):873-4. 876-7. [PubMed ID: [6899769](#)]. [https://doi.org/10.1016/s0001-2092\(07\)61464-1](https://doi.org/10.1016/s0001-2092(07)61464-1).
35. Alfredsdottir H, Bjornsdottir K. Nursing and patient safety in the operating room. *J Adv Nurs.* 2008;**61**(1):29-37. [PubMed ID: [18173734](#)]. <https://doi.org/10.1111/j.1365-2648.2007.04462.x>.
36. Guedon AC, Wauben LS, Overvelde M, Blok JH, van der Elst M, Dankelman J, et al. Safety status system for operating room devices. *Technol Health Care.* 2014;**22**(6):795-803. [PubMed ID: [25159999](#)]. <https://doi.org/10.3233/THC-140854>.
37. Frosini F, Miniati R, Avezzano P, Cecconi G, Dori F, Gentili GB, et al. Development of a web based monitoring system for safety and activity analysis in operating theatres. *Technol Health Care.* 2016;**24**(1):99-109. [PubMed ID: [26409560](#)]. <https://doi.org/10.3233/THC-151073>.
38. Harding K. Risk management in obstetrics. *Obstet Gynaecol Reprod Med.* 2012;**22**(1):1-6. <https://doi.org/10.1016/j.ogrm.2011.10.006>.
39. Biparva AJ, Nikjoo R, Janati A, Arab M, Ostadi A. The implementation framework of active risk management in the operating rooms of hospitals in the country: A qualitative study in Iran. Preprint. *Research Square.* Posted online 12 November 2022. <https://doi.org/10.21203/rs.3.rs-2207573/v1>.
40. Klein VR. Risk Management in Obstetrics and Gynecology. *Clin Obstet Gynecol.* 2019;**62**(3):550-9. [PubMed ID: [31259840](#)]. <https://doi.org/10.1097/GRF.0000000000000473>.
41. Reis CT, Paiva SG, Sousa P. The patient safety culture: a systematic review by characteristics of Hospital Survey on Patient Safety Culture dimensions. *Int J Qual Health Care.* 2018;**30**(9):660-77. [PubMed ID: [29788273](#)]. <https://doi.org/10.1093/intqhc/mzy080>.
42. Marsteller JA, Wen M, Hsu YJ, Bauer LC, Schwann NM, Young CJ, et al. Safety Culture in Cardiac Surgical Teams: Data From Five Programs and National Surgical Comparison. *Ann Thorac Surg.* 2015;**100**(6):2182-9. [PubMed ID: [26330011](#)]. <https://doi.org/10.1016/j.athoracsur.2015.05.109>.
43. Glendon AI, Clarke S, McKenna E. *Human Safety and Risk Management.* 2nd ed. Boca Raton, USA: CRC Press; 2016. <https://doi.org/10.1201/9781420004687>.
44. Chen L. The risk management of medical device-related pressure ulcers based on the Australian/New Zealand Standard. *J Int Med Res.* 2018;**46**(10):4129-39. [PubMed ID: [30092729](#)]. [PubMed Central ID: [PMC6166323](#)]. <https://doi.org/10.1177/0300060518786902>.
45. Nagpal K, Vats A, Ahmed K, Smith AB, Sevdalis N, Jonannsson H, et al. A systematic quantitative assessment of risks associated with poor communication in surgical care. *Arch Surg.* 2010;**145**(6):582-8. [PubMed ID: [20566980](#)]. <https://doi.org/10.1001/archsurg.2010.105>.
46. Puraghaei M, Tabrizi JS, Aslan Abay S, Moharamzadeh P, Ghiami R, Elmdust N. [Investigating the factors and risks in the emergency room of Imam Reza Hospital (AS) in Tabriz]. *Medical journal of mashhad university of medical sciences.* 2015;**58**(6):302-9. Persian. <https://doi.org/10.22038/MJMS.2015.4998>.
47. Kabiri N, Jannati A, Vahed N, Oskouei MM. 109: Risk Management and Patient Safety in Operating Room: A Systematic Review. *BMJ Open.* 2017;**7**(Suppl 1).
48. Aven T. Risk assessment and risk management: Review of recent advances on their foundation. *Eur J Oper Res.* 2016;**253**(1):1-13. <https://doi.org/10.1016/j.ejor.2015.12.023>.
49. Aouicha W, Tlili MA, Sahli J, Mtraoui A, Ajmi T, Said Latiri H, et al. Patient safety culture as perceived by operating room professionals: a mixed-methods study. *BMC Health Serv Res.* 2022;**22**(1):799. [PubMed ID: [35725613](#)]. [PubMed Central ID: [PMC9210674](#)]. <https://doi.org/10.1186/s12913-022-08175-z>.
50. World Health Organization. *Patient safety: Making health care safer.* Geneva, Switzerland: World Health Organization; 2017. 20 p p.
51. Ingvarsdottir E, Halldorsdottir S. Enhancing patient safety in the operating theatre: from the perspective of experienced operating theatre nurses. *Scand J Caring Sci.* 2018;**32**(2):951-60. [PubMed ID: [28940247](#)]. <https://doi.org/10.1111/scs.12532>.
52. Sandelin A, Gustafsson BA. Operating theatre nurses' experiences of teamwork for safe surgery. *Nord J Nurs Res.* 2015;**35**(3):179-85. <https://doi.org/10.1177/0107408315591337>.
53. Stucky CH, Wolf JM. Fire in the Operating Room: Surgical Case Report From a Forced-Air Warming Device Equipment Fire. *J Perianesth Nurs.* 2022;**37**(6):766-9. [PubMed ID: [35811224](#)]. <https://doi.org/10.1016/j.jopan.2022.01.012>.
54. Dalloio L, Raggi A, Sanna T, Mazzetti M, Orsi A, Zanni A, et al. Surveillance of Environmental and Procedural Measures of Infection Control in the Operating Theatre Setting. *Int J Environ Res Public Health.* 2017;**15**(1). [PubMed ID: [29283367](#)]. [PubMed Central ID: [PMC5800145](#)]. <https://doi.org/10.3390/ijerph15010046>.
55. Alshyyab MA, Albsoul R, Fitzgerald G. Factors influencing patient safety culture in operating room in a teaching hospital in Jordan: a qualitative descriptive study. *TQM J.* 2022. <https://doi.org/10.1108/tqm-04-2022-0133>.
56. Omotayo FO. Knowledge Management as an important tool in Organisational Management: A Review of Literature. *Libr Philos Pract.* 2015;**1**(2015):1-23.
57. Institute of Risk Management. *A Risk Practitioners Guide to ISO 31000: 2018.* London, England: Institute of Risk Management; 2018.
58. Helmreich RL, Schaefer H. Team Performance in the Operating Room. In: Bogner MS, editor. *Human Error in Medicine.* Boca Raton, USA: CRC Press; 2018. p. 225-54. <https://doi.org/10.1201/9780203751725-12>.
59. DiCuccio MH. The Relationship Between Patient Safety Culture and Patient Outcomes: A Systematic Review. *J Patient Saf.* 2015;**11**(3):135-42. [PubMed ID: [24583952](#)]. <https://doi.org/10.1097/PTS.000000000000058>.

Table 1. Objectives of the Risk Management Program in the Operating Rooms of Hospitals in Different Countries

No.	Author Name and Year of Publication	Country	Error Type	Risk Management Model	Objectives of the Risk Management Program							
					Quality Improvement	Safety Promotion	Reduction of Costs	Increasing Satisfaction	Prevention and Reduction of Risks	Protecting Financial Losses	Rooting the Risks	
1	Sayed et al., 2013 (20)	Egypt	Any kind of error	WHO safety guidelines	*	*	*	*	*	*	*	*
2	Guo, 2015 (17)	China	Any kind of error	AS-NZS4360 risk management standards	*	*	*	*	*	*	*	*
3	van Beuzekom et al., 2012 (21)	Netherlands	Any kind of error	The Leiden Operating Theatre and Intensive Care Safety (LOTICS) scale	*	*	*	*	*	*	*	*
4	Schimpff, 2007 (22)	USA	Any kind of error	Root-cause analyses	*	*	*	*	*	*	*	*
5	Borie et al., 2018 (23)	France	Fire and burns	Root-cause analyses	*	*	*	*	*	*	*	*
6	Frabetti et al., 2005 (24)	Italy	Microbiological risk	Root-cause analyses	*	*	*	*	*	*	*	*
7	Anghar et al., 2017 (25)	Morocco	Surgical infection, surgical foreign body, medication error, equipment failure, operation error, patient injury, wrong location	The Fuzzy Bayesian Network Model	*	*	*	*	*	*	*	*
8	Silen-Jipponen et al., 2005 (26)	Finland, the USA, and the UK	Human errors	Failure mode and effect analysis	*	*	*	*	*	*	*	*
9	Pomerantz et al., 2011 (27)	USA	Risks associated with skin surgery patients	Failure mode and effect analysis	*	*	*	*	*	*	*	*

10	DeRostier et al., 2019 (18)	USA	Errors related to the process of selection and implantation of intraocular lens (IOL)	Healthcare failure mode and effect analysis	*	*	*
11	Kavosi et al., 2017 (28)	Iran	Any kind of error	Failure mode and effect analysis	*	*	*
12	Le Bourg and Gallois, 2011 (29)	French	Any kind of error	-	*	*	*
13	Lepanluoma et al., 2014 (30)	Finland	Errors related to the neurosurgery patients	WHO surgical safety guidelines	*	*	*
14	American College of Obstetricians and Gynecologists, 2010 (31)	USA	Any kind of error	Universal protocol for preventing wrong site, wrong procedure, and wrong person surgery - WHO surgical safety guidelines	*	*	*
15	American Society of Anesthesiologists Task Force on Operating Room Fires et al., 2008 (32)	USA	Operating room fire	Practice Advisory for the Prevention and Management of Operating Room Fires	*	*	*
16	Bower, 2002 (33)	USA	Any kind of error	Another department's successful program as a model	*	*	*
17	McLain, 1980 (34)	USA	Medical mismanagement, diagnostic and treatment problems, inadequate procedures, equipment failure	Liability control systems	*	*	*

18	Alfredsdottir and Bjornsdottir, 2008 (35)	Iceland	Any kind of error	Root-cause analyses	*	*
19	Tabibzadeh and Jahangiri, 2018 (14)	USA	Any kind of error	Rasmussen's framework	*	*
20	Guedon et al., 2014 (36)	Netherlands	Equipment accidents	Safety status system for operating room devices	*	*
21	Frosini et al., 2016 (37)	Italy	Any kind of error	Web-based monitoring system	*	*

Table 3. Steps for Implementing Risk Management in the Operating Rooms of Hospitals in Different Countries

No.	Author Name and Year of Publication	Steps for Implementing Risk Management	1- Preoperative Evaluation (Log in) 2- Evaluation During Surgery 3- Postoperative Evaluation (Logout) (Model 3)	1- Checking Equipment, Physical Conditions, Environment, Policies, and Procedures to Expose Hazardous Conditions 2- Emphasizing the Safe Operation of Equipment and Preventive Maintenance Plan 3- Early Notification of Preventive Plan Before the Crisis 4- Creating Reporting Channels, Reporting Format, Review, and Timely Documentation and Timely Routing of Error Reports 5- Extensive Internal Training for All Operating Room Staff 6- Finally, Consultation with Operating Room Surgeons (Model 4)
1	Sayed et al., 2003 (20)			*
2	Guo, 2015 (17)			*
3	van Beuzekom et al, 2012 (21)			*
4	Schimpff, 2007 (22)			*
5	Borie et al., 2018 (23)			*
6	Frabetti et al., 2005 (24)			*
7	Amghar et al., 2017 (25)			*
8	Silen-Lipponen et al., 2005 (26)			*
9	Pomerantz et al., 2011 (27)			*
10	DeRosier et al., 2019 (18)		*	
11	Kavosi et al., 2017 (28)		*	
12	Le Bourg and Gallois, 2011 (29)			*
13	Lepanluoma et al., 2013 (30)			*
14	American College of Obstetricians and Gynecologists, 2010 (31)			*

15 American Society of Anesthesiologists Task Force on Operating Room Fires et al., 2008 (32) *

16 Bower, 2002 (33) *

17 McLain, 1980 (34) *

18 Alfredsdottir and Bjornsdottir, 2007 (35) *

19 Tabrizadeh and Jahangiri, 2018 (14) *

20 Guedon et al., 2014 (36) *

21 Frosini et al., 2016 (37) *